

COMPTEL OBSERVATIONS OF THE BLAZAR PKS 1622-297 DURING A GAMMA-RAY HIGH STATE IN 1995

S. Zhang^{1,6}, W. Collmar¹, V. Schönfelder¹, H. Bloemen², W. Hermsen²,
J. Ryan³, K. Bennett⁴, O.R. Williams⁴, and O. Reimer⁵

¹Max-Planck-Institut für extraterrestrische Physik, Garching, Germany

²Space Research Organization Netherlands, Utrecht, The Netherlands

³Space Science Center, University of New Hampshire, Durham, USA

⁴Astrophysics Division, ESTEC, Noordwijk, The Netherlands

⁵NASA/Goddard Space Flight Center, Greenbelt, MD 20771, USA

⁶High Energy Astrophysics Lab, IHEP, P.O.Box 918-3, Beijing, China

ABSTRACT

PKS 1622-297 was detected as a source of γ -rays by the EGRET experiment at energies above 100 MeV during a γ -ray outburst June and July 1995. We analyzed the COMPTEL data (0.75-30 MeV) of this time period to investigate the behaviour of PKS 1622-297 at lower γ -ray energies. The blazar is significantly (5.7σ) detected by COMPTEL at energies above 10 MeV. Below 10 MeV the source is only marginally (3-10 MeV band) or not (below 3 MeV) detected. The summed MeV spectrum shows a 'hard' ($\alpha_{ph} < 2$, $\sim E^{-\alpha}$) shape and, if combined with the simultaneous EGRET spectrum, a spectral break at MeV energies is indicated. We present the COMPTEL results (light curves, spectra) and compare them to results derived in neighbouring energy bands, in particular to the EGRET one.

Key words: γ -rays: observations - galaxies: active - galaxies: quasars: individual: PKS 1622-297.

1. INTRODUCTION

The EGRET experiment aboard CGRO has identified more than 60 blazar-type AGN emitting at high-energy (>100 MeV) γ -rays (Hartman et al. 1999). Most of them show a time-variable γ -ray flux and a few even strong γ -ray flares. One such example is PKS 1622-297, which was detected by EGRET during a γ -ray high state in June and July 1995, and on top of that showed a strong γ -ray flare lasting for a few days (Mattox et al. 1997). After the γ -ray activity of PKS 1622-297 was recognized by EGRET a target-of-opportunity (ToO) observation for CGRO was scheduled which led to the detection of PKS 1622-297 also at hard X- to soft γ -rays by the OSSE experiment (Kurfess et al. 1995).

The COMPTEL experiment (Schönfelder et al. 1993), measuring in the MeV band between 0.75 and 30 MeV, has detected 9 blazars (e.g., Collmar et al. 1999). Among them is PKS 1622-297 which was detected during this γ -ray high state period in 1995. Preliminary MeV results have been reported by Collmar et al. (1997). In this paper we briefly present first results of a detailed analysis of the COMPTEL data of this period, and compare them to results in neighbouring energy bands, in particular to the EGRET band. A more detailed presentation of the results of the analysis will be given in Zhang et al. (2001).

Table 1. COMPTEL observations of PKS 1622-297 during its 4-week γ -ray high state in 1995. The CGRO VPs and their time periods are given. The offset angle refers to the angle between the pointing direction of the instrument and the source position.

VP	Date	Offset angle
421.0	06/06/95-13/06/95	14.5°
422.0	13/06/95-20/06/95	15.2°
423.0	20/06/95-30/06/95	19.2°
423.5	30/06/95-10/07/95	3.0°

2. OBSERVATION AND DATA ANALYSIS

PKS 1622-297 was observed during a four-week observations towards the Galactic Center region from June 6 to July 10, 1995 covering the CGRO viewing periods (VPs) 421 to 423.5. For more details on the observations see Tab. 1.

The analyses were carried out using the standard COMPTEL maximum-likelihood analysis procedure




Figure 1. COMPTEL 10-30 MeV map for the 4-week γ -ray high state of PKS 1622-297 ($\star$) in 1995. The contour lines start at a detection significance level of 3σ with steps of 0.5σ .

(e.g., de Boer et al. 1992) including a filtering technique for background generation (e.g., Bloemen et al. 1994) to derive detection significances, fluxes and flux errors of the γ -ray source in the four standard COMPTEL energy bands. Diffuse emission models have been included in the analysis to subtract off the diffuse galactic γ -ray emission.

3. RESULTS

3.1. Detections

In the sum of the 4 weeks, PKS 1622-297 is detected with a significance of $\sim 5.7\sigma$ in the upper ~ 10 - 30 MeV – COMPTEL band (Fig. 1). Below 10 MeV the source becomes weak. In the 3 - 10 MeV band only a hint is obtained, and below 3 MeV the source is not detected anymore. Above 10 MeV there is evidence for the source in all 4 individual VPs showing that PKS 1622-297 was emitting MeV γ -rays for the whole period. However, the detection significances with about 2 to 3σ each, become weak. The relevant fluxes and upper limits are given in Tab. 2.

3.2. Time Variability

The EGRET (>100 MeV; from Hartman et al. 1999) and COMPTEL 10-30 MeV light curve of the 4 individual CGRO VPs show neither for COMPTEL nor for EGRET strong evidence for time variability (Fig. 2). However, on shorter time scales (~ 1 day) Mattox et al. (1997) reports strong time variability




Figure 2. Light curves for PKS 1622-297 as observed by EGRET at energies > 100 MeV (a) and by COMPTEL in the energy range 10-30 MeV (b) during the 4 individual VPs of the high state in 1995. The EGRET data are from Hartman et al. (1999). The error bars are 1σ .

including an major flare event on top of the γ -ray high state, which occurred within VP 423 and lasted for about 2 days (Fig. 3). We searched the COMPTEL data for a possible simultaneous MeV flare, by subdividing VP 423 according to the EGRET light curve in 3 parts: a pre-flare (TJD 9888.8 - 9892.7), on-flare (TJD 9892.7 - 9894.7), and post-flare part (TJD 9894.7 - 9898.6). The fluxes for the on-flare and post-flare part are higher than the other 10-30 MeV fluxes indicating the presence of the flare also at MeV energies (Fig. 3). We note, however, that the highest COMPTEL flux is found after the flare in the EGRET band had already vanished. This

Table 2. Fluxes and upper limits for PKS 1622-297 for flare state (VPs 421.0-423.5) in units of 10^{-5} ph cm^{-2} s^{-1} . The energy bands are given in MeV. The errors are 1σ and the upper limits are 2σ . An upper limit is given when the significance of an individual flux value is less than 1σ .

Period (VP)	0.75-1	1-3	3-10	10-30
421.0	<28.3	<25.2	<10.7	3.4 ± 1.7
422.0	18.3 ± 13.6	<28.0	<8.9	3.2 ± 1.5
423.0	18.1 ± 12.5	<23.9	<10.0	5.1 ± 1.5
423.5	<21.5	<20.8	<9.7	1.8 ± 1.0
421.0-423.5	<17.4	<11.2	2.0 ± 1.9	3.5 ± 0.7




Figure 3. High time resolution EGRET light curve for PKS 1622-297 (a) and compared to COMPTEL light curves (b, c), where the major flaring period is better resolved in time. Fig. 3c is the same as Fig. 3b with the two high fluxes combined to show the enhanced MeV emission around the time of EGRET major flare. The EGRET data are from Mattox et al. (1997).

indicates a possible time delay of a few days between the two bands in the sense that the higher energies come first.

3.3. Energy Spectra

We have added the COMPTEL data of the 4-week high-state period to generate an average MeV spectrum. Due to the non-detections below 3 MeV only 2 spectral points are derived, which, nevertheless, indicate a hard ($\alpha_{ph} < 2$, $\sim E^{-\alpha}$) spectrum of PKS 1622-297 at MeV energies (Fig. 4). We also generated the simultaneous EGRET spectrum of PKS 1622-297, which is between 30 MeV and 10 GeV sufficiently fitted by a power-law shape with a photon index of ~ 2.2 . If the simultaneous EGRET and COMPTEL spectra are combined (Fig. 4), a spectral break at MeV energies becomes obvious. A fit of this simultaneous EGRET and COMPTEL spectrum with a broken power-law model indicates the break energy to be around 20 MeV. Due to the marginal COMPTEL detections below 10 MeV, the amount of the spectral break, i.e. the change in photon index from the higher to the lower energies, is not well determined. Its lower limit, however, should be larger than 0.6.




Figure 4. The average COMPTEL spectrum (a) and the combined COMPTEL/EGRET spectrum (b) together with the best-fitting broken power-law model for the 4-week γ -ray high-state period are shown. The evidence for the spectral break around ~ 20 MeV is obvious.

After the γ -ray activity of PKS 1622-297 had been recognized by EGRET, a ToO observation (VP 423.5) was carried out, where OSSE participated. This provided the possibility to measure the spectrum down to 0.05 MeV. Mattox et al. (1997) reported that the OSSE and EGRET data itself can be represented by power-law shapes with similar photon indices of 2.0 ± 0.2 and 2.2 ± 0.1 , respectively (Fig. 5). However, fitting a simple power-law shape to the combined OSSE/EGRET data, they obtained a photon spectral index of 1.87 ± 0.02 . We have analysed the COMPTEL data of VP 423.5 and have added the spectral results to this combined OSSE-EGRET spectrum (Fig. 5). Although these COMPTEL data are not really constraining, they – at first glance – support a more complex spectral shape at γ -ray energies than the simple power-law model.

4. DISCUSSION

In general, PKS 1622-297 shows a similar behaviour as is observed in other such sources, like PKS 0528+134 and 3C 279 for example. During flaring events observed by EGRET, PKS 1622-297 is detected – predominantly – at the upper COMPTEL energies, resulting in a 'hard' MeV spectrum and showing a spectral break in the combined COMPTEL-EGRET spectrum. These facts indicate




Figure 5. The combined simultaneous OSSE/COMPTEL/EGRET spectrum for the ToO period (VP 423.5). The EGRET and OSSE data and the relevant fit lines are from Mattox et al. (1997). The dotted line symbolizes the OSSE fit, the dashed-dotted line the EGRET fit, and the solid line the combined EGRET/OSSE fit.

that the flare is more efficient at higher γ -ray energies. Such a behaviour can generally be explained in a multi-component γ -ray emission scenario, consisting of synchrotron-self Compton and external Comptonization components. Because they have different dependencies on the bulk Lorentz factor, a change in bulk Lorentz factor could result in increased emission at mainly the higher-energy γ -rays (e.g., Böttcher & Collmar 1998).

On top of the already high γ -ray level, PKS 1622-297 showed a major flare which lasted for a few days. COMPTEL measures enhanced γ -ray emission during this period. This suggests that the flare is also visible at MeV energies, however with reduced amplitude. An intriguing feature of this flare is the possible time delay of a few days between the γ -rays above 100 MeV and the 10-30 MeV band, indicating that the opacity of the γ -ray emitting region might be different for the different energy bands. Such a delay has not yet been observed in another blazar. A measurement of this delay, together with the ratio of the flare amplitudes in the different bands, has the potential to improve our insights on the γ -ray emitting region.

5. SUMMARY

During a 4-week γ -ray activity period observed at energies above 100 MeV by EGRET, PKS 1622-297 is also significantly detected at lower-energy γ -rays (>10 MeV) by COMPTEL. This detection together with the marginal detection and non-detection at lower COMPTEL energies indicates a 'hard' ($\alpha_{ph} < 2$, $\sim E^{-\alpha}$) MeV spectrum. The combined COMPTEL-EGRET spectrum requires a spectral break around ~ 20 MeV.

On top of this γ -ray high state, EGRET observed a major flare which lasted for about 2 days. COMPTEL observes enhanced MeV emission (10 - 30 MeV band) during that time. However, the COMPTEL flux is still high when the EGRET one is already down, indicating a possible time delay of a few days between the two bands.

ACKNOWLEDGEMENTS

This research was supported by the German government through DLR grant 50 QV 9096 8, by NASA under contract NAS5-26645, and by the Netherlands Organisation for Scientific Research NWO.

REFERENCES

- Bloemen H., Hermsen W., Swanenburg B.N., et al., 1994, ApJS 92, 419
- de Boer, H., Bennett K., Bloemen H., et al., 1992, In: *Data Analysis in Astronomy IV*, eds. V. Di Gesu et al., New York, USA, p. 241
- Böttcher M., Collmar W., 1998, A&A 329, 57
- Collmar W., Bennett K., Bloemen H., et al., 1997, in *Proceedings of the Fourth Compton Symposium*, eds. C.D. Dermer, M.S. Strickman, J.D. Kurfess (New York: AIP Conf. Proc. 410), p. 1341
- Collmar W., Bennett K., Bloemen H., et al., 1999, *Astrophys. Lett. & Communications*, Vol. 39, p. 57
- Hartman R.C., Bertsch D.L., Bloom S.D., et al., 1999, ApJS 123, 79
- Kurfess J.D., Grove J.E., McNaron-Brown K., et al., 1995, IAU circular 6185
- Mattox J.R., Wagner S.J., Malkan M., et al., 1997, ApJ 476, 692
- Schönfelder V., Aarts H., Bennett K., et al., 1993, ApJS 86, 657
- Zhang S., Collmar W., Bennett K., et al., 2001, in prep.